

STRATIFIED BUNDLES AND ÉTALE FUNDAMENTAL GROUP

HÉLÈNE ESNAULT AND XIAOTAO SUN

ABSTRACT. On X projective smooth over an algebraically closed field of characteristic $p > 0$, we show that all irreducible stratified bundles have rank 1 if and only if the commutator $[\pi_1, \pi_1]$ of the étale fundamental group π_1 is a pro- p -group, and we show that the category of stratified bundles is semi-simple with irreducible objects of rank 1 if and only if π_1 is abelian without p -power quotient. This answers positively a conjecture by Gieseker [4, p. 8].

1. INTRODUCTION

Let X be a smooth projective variety defined over an algebraically closed field k of characteristic $p > 0$. In [4], stratified bundles are defined and studied. It is shown that they are a characteristic $p > 0$ analog to complex local systems over smooth complex algebraic varieties. In particular, Gieseker shows [4, Theorem 1.10]

- (i) If every stratified bundle is trivial, then π_1 is trivial.
- (ii) If all the irreducible stratified bundles have rank 1, then $[\pi_1, \pi_1]$ is a pro- p -group.
- (iii) If every stratified bundle is a direct sum of stratified line bundles, then π_1 is abelian without non-trivial p -power quotient.

Here π_1 is the étale fundamental group based at some geometric point. He conjectures that in the three statements, the “if” can be replaced by “if and only if”. The aim of this note is to give a positive answer to Gieseker’s conjecture (see Theorem 3.9).

The converse to (i) is the main theorem of [3], and is analog to Malcev-Grothendieck theorem ([10], [5]) asserting that if the étale fundamental group of a smooth complex projective variety is trivial, then there is no non-trivial bundle with a flat connection.

Date: December 20, 2011.

The first author is supported by the SFB/TR45 and the ERC Advanced Grant 226257. The second author is supported by NBRPC 2011CB302400, NSFC60821002/F02 and NSFC 10731030.

The complex analog to the converse to (iii) relies on the following.

If X is an abelian variety over a field k , Mumford [11, Section 16] showed that a non-trivial line bundle L which is algebraically equivalent to 0 fulfills $H^i(X, L) = 0$ for all $i \geq 0$. If $k = \mathbb{C}$, the field of complex numbers, L carries a unitary flat connection with underlying local system ℓ , and $H^1(X, L) = 0$ implies that $H^1(X, \ell) = 0$. In fact, more generally, if X is any complex manifold with abelian topological fundamental group, and ℓ is a non-trivial rank 1 local system, then $H^1(X, \ell) = 0$ (see Remark 3.11).

Finally the complex analog to the converse to (ii) and (iii) together says that if X is a smooth complex variety, then all irreducible complex local systems on X have rank 1 if and only if π_1 is abelian. However, the semi-simplicity statement has a different phrasing: all irreducible complex local systems on X have rank 1 and the category is semi-simple if and only if π_1 is a finite abelian group (see Claim 3.13 for a precise statement).

The proof of the converse to (ii) is done in Section 2. It relies on a consequence of the proof of the main theorem in [3], which is formulated in Theorem 2.3. It is a replacement for the finite generation of the topological fundamental group in complex geometry (see the analog statement in Remark 3.12). The proof of the converse to (iii) is done in Section 3. The key point to show the converse to (iii) is Theorem 3.4. Its proof relies on the construction (Proposition 3.5) of a moduli space of non-trivial extensions, with a rational map f induced by Frobenius pullback, which allows us to use a theorem of Hrushovski.

Acknowledgements: We thank the department of Mathematics of Harvard University for its hospitality during the preparation of this note. We thank Nguyễn Duy Tân for a very careful reading of a first version, which allowed us to improve the redaction. We thank the referee of the first version who pointed out a mistake, corrected in Section 3. The second named author would like to thank his colleague Nanhua Xi for discussions related to Claim 3.3.

2. STRATIFIED BUNDLES

Let k be an algebraically closed field of characteristic $p > 0$, and X a smooth connected projective variety over k . A stratified bundle on X is by definition a coherent \mathcal{O}_X -module \mathcal{E} with a homomorphism

$$\nabla : \mathcal{D}_X \rightarrow \mathcal{E}nd_k(\mathcal{E})$$

of \mathcal{O}_X -algebras, where \mathcal{D}_X is the sheaf of differential operators acting on the structure sheaf of X . By a theorem of Katz (cf. [4, Theorem

1.3]), it is equivalent to the following definition. (The terminology is not unique: stratified bundles from the following definition are called flat bundles in [4], F -divided sheaves in [2]).

Definition 2.1. A stratified bundle on X is a sequence of bundles

$$E = \{E_0, E_1, E_2, \dots, \sigma_0, \sigma_1, \dots\} = \{E_i, \sigma_i\}_{i \in \mathbb{N}}$$

where $\sigma_i : F_X^* E_{i+1} \rightarrow E_i$ is a \mathcal{O}_X -linear isomorphism, and $F_X : X \rightarrow X$ is the absolute Frobenius.

A morphism $\alpha = \{\alpha_i\} : \{E_i, \sigma_i\} \rightarrow \{F_i, \tau_i\}$ between two stratified bundles is a sequence of morphisms $\alpha_i : E_i \rightarrow F_i$ of \mathcal{O}_X -modules such that

$$\begin{array}{ccc} F_X^* E_{i+1} & \xrightarrow{F_X^* \alpha_{i+1}} & F_X^* F_{i+1} \\ \sigma_i \downarrow & & \downarrow \tau_i \\ E_i & \xrightarrow{\alpha_i} & F_i \end{array}$$

is commutative. The category $\mathbf{str}(X)$ of stratified bundles is abelian, rigid, monoidal. To see it is k -linear, it is better to define stratified bundles and morphisms in the relative version: objects consist of

$$E' = \{E_0 = E'_0, E'_1, E'_2, \dots, \sigma'_0, \sigma'_1, \dots\} = \{E'_i, \sigma'_i\}_{i \in \mathbb{N}}$$

where E'_i is a bundle on the i -th Frobenius twist $X^{(i)}$ of X ,

$$\sigma'_i : F_{i,i+1}^* E'_{i+1} \rightarrow E'_i$$

is a $\mathcal{O}_{X^{(i)}}$ -linear isomorphism, and $F_{i,i+1} : X^{(i)} \rightarrow X^{(i+1)}$ is the relative Frobenius, the morphisms are the obvious ones. A rational point $a \in X(k)$ yields a fiber functor $\omega_a : \mathbf{str}(X) \rightarrow \mathbf{vec}_k$, $E \mapsto a^* E_0$ with values in the category of finite dimensional vector spaces. Thus $(\mathbf{str}(X), \omega_a)$ is a Tannaka category ([2, Section 2.2]), and one has an equivalence of categories

$$(2.1) \quad \mathbf{str}(X) \xrightarrow{\omega_a \cong} \mathbf{rep}_k(\pi^{\mathbf{str}})$$

where

$$\pi^{\mathbf{str}} = \mathrm{Aut}^{\otimes}(\omega_a)$$

is the Tannaka group scheme, and $\mathbf{rep}_k(\pi^{\mathbf{str}})$ is the category of finite dimensional k -representations of $\pi^{\mathbf{str}}$. Let $\pi_1 := \pi_1^{\acute{\mathrm{e}}\mathrm{t}}(X, a)$ be the étale fundamental group of X . In [4, Theorem 1.10], D. Gieseker proved the following theorem.

Theorem 2.2. (*Gieseker*): *Let X be a smooth projective variety over an algebraically closed field k , then*

- (i) *If every stratified bundle is trivial, then π_1 is trivial.*

- (ii) *If all the irreducible stratified bundles have rank 1, then $[\pi_1, \pi_1]$ is a pro- p -group.*
- (iii) *If every stratified bundle is a direct sum of stratified line bundles, then π_1 is abelian with no p -power order quotient.*

Then Gieseker conjectured that the converse of the above statements might be true. In [3], it is proven that the converse of statement (i) is true. The aim of this section is to prove the converse of (ii). We prove the converse of (iii) in Section 3.

The proof relies on the following theorem extracted from [3], and which plays a similar rôle as the finite generation of the topological fundamental group π_1^{top} in complex geometry (see Remark 3.12).

Fixing an ample line bundle $\mathcal{O}_X(1)$ on X , recall that E is said to be μ -stable if for all coherent subsheaves $U \subset E$ one has $\mu(U) < \mu(E)$, where $\mu(E)$ is the slope which is defined to be the degree of E with respect to $\mathcal{O}_X(1)$ divided by the rank of E .

Theorem 2.3. *Let X be a smooth projective variety over an algebraically closed field k . If there is a stratified bundle $E = (E_n, \sigma_n)_{n \in \mathbb{N}}$ of rank $r \geq 2$ on X , where $\{E_n\}_{n \in \mathbb{N}}$ are μ -stable, then there exists an irreducible representation $\rho : \pi_1 \rightarrow GL(V)$ with finite monodromy, where V is a r -dimensional vector space over \mathbb{F}_p .*

Proof. By the proof of [3, Theorem 3.15], there is a smooth projective model $X_S \rightarrow S$ of $X \rightarrow \text{Spec } k$, with model $a_S \in X_S(S)$ of $a \in X(k)$, where S is smooth affine over \mathbb{F}_p , together with a quasi-projective model $M_S \rightarrow S$ of the moduli of stable vector bundles considered in [3, Section 3], there is a closed point $u \rightarrow M_S$ of residue field \mathbb{F}_q , corresponding to a bundle \mathcal{E} over X_s , stable over $X_{\bar{s}} = X_s \otimes \bar{\mathbb{F}}_p$, where s is the closed point u viewed as a closed point of S , such that $(F^m)^*\mathcal{E} \cong \mathcal{E}$, for some $m \in \mathbb{N} \setminus \{0\}$, where F is the Frobenius of X_s/\mathbb{F}_q . Thus, as explained in [3, Theorem 3.15], \mathcal{E} trivializes over a Lang torsor $Y \rightarrow X_s$ [9, Satz 1.4], thus its base change $\mathcal{E}_{\bar{s}}$ to $X_{\bar{s}}$ trivializes over $Y \times_s \bar{s} \rightarrow X_{\bar{s}}$. This trivialization defines an irreducible representation $\rho' : \pi_1^{\text{ét}}(X_{\bar{s}}, b) \rightarrow GL(\mathcal{E}_b)$, where $b = a_S \otimes \bar{s}$. As the specialization homomorphism $\text{sp} : \pi_1 \rightarrow \pi_1^{\text{ét}}(X_{\bar{s}}, b)$ is surjective ([13, Exposé X, Théorème 3.8]), the composite $\rho = \text{sp} \circ \rho'$ is a representation of π_1 with the same finite irreducible monodromy. This is a solution to the problem. □

We now use the following two elementary lemmas.

Lemma 2.4 (See Chapter 8, Proposition 26 of [12]). *Let G be a finite p -group and $\rho : G \rightarrow GL(V)$ be a representation on a vector space*

$V \neq 0$ over a field k of characteristic p . Then

$$V^G = \{ v \in V \mid \rho(g)v = v, \forall g \in G \} \neq 0.$$

Lemma 2.5. *Let G be a finite group and $\rho : G \rightarrow GL(V)$ be an irreducible representation on a finite dimensional vector space V over an algebraically closed field k of characteristic $p > 0$. If the commutator $[G, G]$ of G is a p -group, then $\dim(V) = 1$.*

Proof. If $[G, G]$ is a p -group, by Lemma 2.4, there exists a $0 \neq v \in V$ such that $\rho(g_1)\rho(g_2)v = \rho(g_2)\rho(g_1)v$ for any $g_1, g_2 \in G$. Since V is an irreducible G -module, the sub-vector space spanned by the orbit $\rho(G)v$ is V itself. Thus for all $w \in V$, all $g \in G$, there are $a_g(w) \in k$, such that

$$w = \sum_{g \in G} a_g(w) \rho(g)v.$$

On the other hand, for all $g_1, g_2, g \in G$, one has $g_1g_2 = g_2g_1c$ for some $c \in [G, G]$, thus $g_1g_2g = g_2g_1cg = g_2g_1gc'$, for some $c' \in [G, G]$. We conclude that $\rho(g_1)\rho(g_2)w = \rho(g_2)\rho(g_1)w$ for any $g_1, g_2 \in G$, that is $\rho(G)$ is abelian. As ρ is irreducible, V must have dimension 1. \square

Theorem 2.6. *Let X be a smooth projective variety over an algebraically closed field k . Then all irreducible stratified bundles on X have rank 1 if and only if the commutator $[\pi_1, \pi_1]$ of π_1 is a pro- p -group.*

Proof. One direction is the (ii) in Theorem 2.2. We prove the converse. Assume that $[\pi_1, \pi_1]$ is a pro- p -group. Let $E = (E_n, \sigma_n)_{n \in \mathbb{N}}$ be an irreducible stratified bundle of rank $r \geq 1$ on X . Assume $r \geq 2$. Then by [3, Proposition 2.3], there is a $n_0 \in \mathbb{N}$ such that the stratified bundle $E(n_0) := (E_n, \sigma_n)_{n \geq n_0}$ is a successive extension of stratified bundles $U = (U_n, \tau_n)$ with underlying bundles U_n being μ -stable. But E being irreducible implies that $E(n_0)$ is irreducible as well. Hence all the E_n are μ -stable for $n \geq n_0$. By Theorem 2.3, there is an irreducible representation $\rho : \pi_1 \rightarrow GL(V)$ of dimension $r \geq 2$ with finite monodromy over an algebraically closed field of characteristic $p > 0$. By Lemma 2.5, this is impossible. Thus $r = 1$. \square

3. EXTENSIONS OF STRATIFIED LINE BUNDLES

In this section, we prove the following theorem.

Theorem 3.1. *Let X be a smooth projective connected variety over an algebraically closed field k of characteristic $p > 0$. If the étale*

fundamental group π_1 of X is abelian and has no non-trivial p -power order quotient, then any extension

$$0 \rightarrow \mathbb{L} \rightarrow \mathbb{V} \rightarrow \mathbb{L}' \rightarrow 0$$

in $\mathbf{str}(X)$ is split when \mathbb{L} and \mathbb{L}' are rank 1 objects.

We start the proof by the following proposition, which can be considered as a generalization of [3, Proposition 2.4].

Proposition 3.2. *Let L be a line bundle on X such that $(F_X^*)^a L$ is isomorphic to L , for some non zero natural number a , where $F_X : X \rightarrow X$ is the absolute Frobenius map. Then*

$$(F_X^*)^a : H^1(X, L) \rightarrow H^1(X, L)$$

is nilpotent if π_1 is abelian without non-trivial p -power quotient.

Proof. We choose an isomorphism $(F_X^*)^a L \cong L$, or equivalently an isomorphism $L^{p^a-1} \cong \mathcal{O}_X$. This define a Kummer cover $\phi : Y \rightarrow X$ with Galois group $H = \mathbb{Z}/(p^a - 1)$ such that

$$\phi^* L = \mathcal{O}_Y.$$

One has an exact sequence (recall that $\pi_1 := \pi_1^{\text{ét}}(X, \bar{a})$)

$$1 \rightarrow \pi_1^{\text{ét}}(Y, \bar{b}) \rightarrow \pi_1 \rightarrow H \rightarrow 1,$$

where \bar{b} is a geometric point of Y above the geometric point \bar{a} of X .

Claim 3.3. $\pi_1^{\text{ét}}(Y, \bar{b})$ is abelian without p -power quotient.

Proof. Since π_1 is commutative, the kernel of any quotient map

$$\pi_1^{\text{ét}}(Y, \bar{b}) \twoheadrightarrow K$$

is normal in π_1 . Taking for K a finite p -group defines the push-out exact sequence

$$1 \rightarrow K \rightarrow G \rightarrow H \rightarrow 1$$

where G is a quotient of π_1 , thus is commutative and has no p -power quotient. Since K splits in G , $K = \{1\}$. \square

Then, by [3, Proposition 2.4], there is an integer $N > 0$ such that

$$(F_Y^*)^N : H^1(Y, \mathcal{O}_Y) \rightarrow H^1(Y, \mathcal{O}_Y)$$

is a zero map, which implies that

$$\phi^* \cdot (F_X^*)^{Na} = (F_Y^*)^{Na} \cdot \phi^* : H^1(X, L) \rightarrow H^1(Y, \mathcal{O}_Y)$$

is a zero map. But $\phi^* : H^1(X, L) \hookrightarrow H^1(Y, \phi^* L) = H^1(Y, \mathcal{O}_Y)$ is injective, thus $(F_X^*)^{Na} : H^1(X, L) \rightarrow H^1(X, L)$ is a zero map. \square

Proof of Theorem 3.1. By twisting with $(\mathbb{L}')^{-1}$, we may assume that $\mathbb{L}' = \mathbb{L}$, the trivial object in $\mathbf{str}(X)$. We prove Theorem 3.1 by contradiction. If there exists a nontrivial extension

$$(3.1) \quad 0 \rightarrow \mathbb{L} \rightarrow \mathbb{V} \rightarrow \mathbb{L} \rightarrow 0$$

in $\mathbf{str}(X)$, then, by definition, there is a set

$$\mathbb{E}_X = \{ (L_i \hookrightarrow V_i \twoheadrightarrow \mathcal{O}_X) \}_{i \in \mathbb{N}}$$

of isomorphism classes of non-trivial extensions on X such that

$$(L_i \hookrightarrow V_i \twoheadrightarrow \mathcal{O}_X) \cong (F_X^* L_{i+1} \hookrightarrow F_X^* V_{i+1} \twoheadrightarrow F_X^* \mathcal{O}_X)$$

for any $i \in \mathbb{N}$ modulo scale. This means that there is $\lambda_i \in k^\times = \Gamma(X, \mathcal{O}_X^\times)$ such that the push down of $(L_i \hookrightarrow V_i \twoheadrightarrow \mathcal{O}_X)$ by $L_i \xrightarrow{\lambda_i} L_i$ is isomorphic as an extension to $(F_X^* L_{i+1} \hookrightarrow F_X^* V_{i+1} \twoheadrightarrow \mathcal{O}_X)$. In particular, L_i is isomorphic as a line bundle to $F_X^*(L_{i+1})$.

If \mathbb{E}_X is a finite set, then there are only finitely many non-isomorphic L_i , thus there is a $a \in \mathbb{N} \setminus \{0\}$ such that $(F_X^*)^a L_i \cong L_i$ for all i . We obtain a contradiction by Proposition 3.2.

If \mathbb{E}_X is an infinite set, by Theorem 3.4 below, there is a nontrivial extension

$$(3.2) \quad 0 \rightarrow L \rightarrow V \rightarrow \mathcal{O}_{X_{\bar{s}}} \rightarrow 0$$

on a good reduction $X_{\bar{s}}$ of X (thus over $\overline{\mathbb{F}}_p$) such that for an integer $a > 0$

$$0 \rightarrow (F_{X_{\bar{s}}}^*)^a L \rightarrow (F_{X_{\bar{s}}}^*)^a V \rightarrow (F_{X_{\bar{s}}}^*)^a \mathcal{O}_{X_{\bar{s}}} \rightarrow 0$$

is isomorphic to the extension (3.2) modulo scale. Then, by using Proposition 3.2 to $X_{\bar{s}}$, we obtain again a contradiction since $\pi_1^{\text{ét}}(X_{\bar{s}}, a_{\bar{s}})$, which is a quotient of π_1 via the specialization map, is also abelian without non-trivial p -power quotient. Here $a_{\bar{s}}$ is a specialization of the geometric point \bar{a} used to define π_1 . \square

Theorem 3.4. *Let X be a smooth projective variety over an algebraically closed field k . If there exists an infinite set \mathbb{E}_X of equivalent classes of nontrivial extensions*

$$(L_i \hookrightarrow V_i \twoheadrightarrow \mathcal{O}_X), \quad i \in \mathbb{N}$$

satisfying $(L_i \hookrightarrow V_i \twoheadrightarrow \mathcal{O}_X) \cong (F_X^ L_{i+1} \hookrightarrow F_X^* V_{i+1} \twoheadrightarrow F_X^* \mathcal{O}_X)$ for any $i \in \mathbb{N}$, modulo scale, then there exists a nontrivial extension*

$$0 \rightarrow L \rightarrow V \rightarrow \mathcal{O}_{X_{\bar{s}}} \rightarrow 0$$

on a good reduction $X_{\bar{s}}$ of X (over $\overline{\mathbb{F}}_p$) such that for some $a > 0$

$$((F_{X_{\bar{s}}}^*)^a L \hookrightarrow (F_{X_{\bar{s}}}^*)^a V \twoheadrightarrow (F_{X_{\bar{s}}}^*)^a \mathcal{O}_{X_{\bar{s}}}) \cong (L \hookrightarrow V \twoheadrightarrow \mathcal{O}_{X_{\bar{s}}})$$

modulo scale.

We now prove Theorem 3.4. Let $X_S \rightarrow S$ be a smooth projective model of X/k , endowed with a section, where S is a smooth affine irreducible variety over \mathbb{F}_q with $H^1(X, \mathcal{O}_{X_S}) = H^0(S, \mathcal{O}_S)$. So $X_S \rightarrow S$ has smooth projective geometrically irreducible fibers. Then we have

Proposition 3.5. *There exists a reduced S -scheme $M \rightarrow S$ of finite type and a rational map $f : M \dashrightarrow M$ over S such that*

- (1) *For any field extension $K \supset \mathbb{F}_q$, the set $M(K) = \{e\}$ of K -valued points consists of isomorphism classes of non-trivial extensions $L \hookrightarrow V \twoheadrightarrow \mathcal{O}_{X_s}$ modulo scale on X_s , where X_s is the fiber of $X_S \rightarrow S$ at the image $s \in S(K)$ of $e \in M(K)$.*
- (2) *The set \mathbb{E}_X of Theorem 3.4 minus finitely many elements lies in $M(k)$.*
- (3) *The rational map $f : M \dashrightarrow M$ is well-defined at $e = (L \hookrightarrow V \twoheadrightarrow \mathcal{O}_{X_{s_0}}) \in M(K)$ if and only if the pull-back extension*

$$0 \rightarrow F_{X_s}^* L \rightarrow F_{X_s}^* V \rightarrow F_{X_s}^* \mathcal{O}_{X_s} \rightarrow 0$$

under the absolute Frobenius F_{X_s} does not split, in which case

$$f(e) = (F_{X_s}^* L \hookrightarrow F_{X_s}^* V \twoheadrightarrow F_{X_s}^* \mathcal{O}_{X_s}) \in M(K).$$

Proof. The line bundles $\{L_i\}_{i \in \mathbb{N}}$ occurring in \mathbb{E}_X must satisfy

$$L_i \cong L_{i+1}^p, \quad \forall i \in \mathbb{N},$$

so they are infinitely p -divisible, thus lie in $\text{Pic}^\tau(X)(k)$, the group of numerically trivial line bundles over k ([8, 9.6]).

We first assume $\mathbb{L} = \mathbb{I}$. We shrink S so that $H^1(\mathcal{O}_{X_S})$ is locally free and commutes with base change. Let $H^1(\mathcal{O}_{X_S})$ denote the first direct image of \mathcal{O}_{X_S} under $X_S \rightarrow S$, then

$$\pi_S : M = \mathbb{P}(H^1(\mathcal{O}_{X_S})^\vee) \rightarrow S$$

is the moduli space of isomorphism classes of non-trivial extensions $\mathcal{O}_{X_s} \hookrightarrow V \twoheadrightarrow \mathcal{O}_{X_s}$. In the diagram

$$\begin{array}{ccccc} & & F_{X_S} & & \\ & \nearrow & & \searrow & \\ X_S & \xrightarrow{F} & X'_S & \longrightarrow & X_S \\ & \searrow & \downarrow & & \downarrow \\ & & S & \xrightarrow{F_S} & S, \end{array}$$

F_S, F_{X_S} are the absolute Frobenius and $F : X_S \rightarrow X'_S$ is the relative Frobenius. Then the "Frobenius pullback" induces

$$(3.3) \quad F^* : F_S^* H^1(\mathcal{O}_{X_S}) = H^1(\mathcal{O}_{X'_S}) \rightarrow H^1(\mathcal{O}_{X_S})$$

which is \mathcal{O}_S -linear and commutes with base change. Let

$$\pi_S^* H^1(\mathcal{O}_{X_S})^\vee \rightarrow \mathcal{O}_M(1) \rightarrow 0$$

be the universal rank one quotient, which, pulled back by the absolute Frobenius $F_M : M \rightarrow M$, defines the \mathcal{O}_M -linear map

$$(3.4) \quad \pi_S^* F_S^* H^1(\mathcal{O}_{X_S})^\vee \rightarrow \mathcal{O}_M(p) \rightarrow 0$$

(since $F_M^* \pi_S^* = \pi_S^* F_S^*$ and $F_M^* \mathcal{O}_M(1) := \mathcal{O}_M(p)$). Combining the dual

$$\pi_S^*(F^{*\vee}) : \pi_S^* H^1(\mathcal{O}_{X_S})^\vee \rightarrow \pi_S^* F_S^* H^1(\mathcal{O}_{X_S})^\vee$$

of (3.3) with (3.4) defines the \mathcal{O}_M -linear map

$$\pi_S^* H^1(\mathcal{O}_{X_S})^\vee \rightarrow \mathcal{O}_M(p).$$

By the universal property, this is the same as a rational map

$$f : M \dashrightarrow M$$

over S . To see that f satisfies (3), let $e \rightarrow M$ be a point with image $s \rightarrow S$ (under $M \xrightarrow{\pi_S} S$), which corresponds a nontrivial extension

$$e = (\mathcal{O}_{X_s} \hookrightarrow V \twoheadrightarrow \mathcal{O}_{X_s})$$

where X_s is the fiber of $X_S \rightarrow S$ at $s \rightarrow S$. Then f is well-defined at $e \rightarrow M$ if and only if the \mathcal{O}_M -linear map

$$\pi_S^* H^1(\mathcal{O}_{X_S})^\vee \rightarrow \mathcal{O}_M(p)$$

at $e \rightarrow M$ is surjective. By definition, this is equivalent to saying that the \mathcal{O}_S -linear map (3.3) at $s \rightarrow S$, which is

$$F^* : F_{k(s)}^* H^1(\mathcal{O}_{X_s}) = H^1(\mathcal{O}_{X'_s}) \rightarrow H^1(\mathcal{O}_{X_s}),$$

is not trivial at $F_{k(s)}^*([e]) \in H^1(\mathcal{O}_{X'_s})$ (i.e. $F^*(F_{k(s)}^*([e])) \neq 0$), where

$$[e] \in H^1(\mathcal{O}_{X_s})$$

denotes the corresponding class of $e = (\mathcal{O}_{X_s} \hookrightarrow V \twoheadrightarrow \mathcal{O}_{X_s})$. Since

$$F_{X_s}^* : H^1(\mathcal{O}_{X_s}) \xrightarrow{F_{k(s)}^*} H^1(\mathcal{O}_{X'_s}) \xrightarrow{F^*} H^1(\mathcal{O}_{X_s}),$$

we have $F_{X_s}^*([e]) = F^*(F_{k(s)}^*([e]))$. Thus f is well-defined at

$$e = (\mathcal{O}_{X_s} \hookrightarrow V \twoheadrightarrow \mathcal{O}_{X_s}) \in M$$

if and only if $(F_{X_s}^* \mathcal{O}_{X_s} \hookrightarrow F_{X_s}^* V \twoheadrightarrow F_{X_s}^* \mathcal{O}_{X_s})$ is not splitting, and

$$f(e) = (F_{X_s}^* \mathcal{O}_{X_s} \hookrightarrow F_{X_s}^* V \twoheadrightarrow F_{X_s}^* \mathcal{O}_{X_s}).$$

The objects $(\mathcal{O}_X \hookrightarrow V_i \twoheadrightarrow \mathcal{O}_X) \in \mathbb{E}_X$ ($i \in \mathbb{N}$) define points $e_i \rightarrow M$ over $s = \text{Spec}(k) \rightarrow S$, and f is well-defined at e_i with $f(e_i) = e_{i-1}$ for $i > 1$. This finishes the proof if $\mathbb{L} = \mathbb{I}$.

If $\mathbb{L} \neq \mathbb{L}$, after removing a finite number of elements in the set \mathbb{E}_X , we can assume that all line bundles $\{L_i\}_{i \in \mathbb{N}}$ occurring in \mathbb{E}_X satisfy

$$H^0(X, L_i) = 0, \quad H^1(L_i) \neq 0 \quad i \in \mathbb{N}.$$

Let $\text{Pic}_{X_S}^\tau \rightarrow S$ be the torsion component of the identity of the Picard scheme [8, 9.6]. Let \mathcal{L} be the universal line bundle on $X_S \times_S \text{Pic}_{X_S}^\tau$. We define

$$\mathcal{N}_0 = \{ t \in \text{Pic}_{X_S}^\tau \mid H^1(\mathcal{L}_t) \neq 0 \}$$

with its reduced structure. By semi-continuity of cohomology, it is a closed sub-scheme of $\text{Pic}_{X_S}^\tau$. If \mathcal{N}_i is defined, let

$$\mathcal{N}_{i+1} = \{ t \in \mathcal{N}_i \mid H^1(\mathcal{L}_t^{p^{i+1}}) \neq 0 \}.$$

By the noetherian property, $\mathcal{N}_0 \supseteq \mathcal{N}_1 \supseteq \cdots \mathcal{N}_i \supseteq \mathcal{N}_{i+1} \supseteq \cdots$ terminates. Then, there is a $k_0 \geq 0$ such that

$$\mathcal{N}_i = \mathcal{N}_{k_0}, \quad \forall i \geq k_0.$$

The line bundles $\{L_{i+p^{k_0}}\}_{i \in \mathbb{N}}$ occurring in $\mathbb{E}_X(p^{k_0})$, where

$$\mathbb{E}_X(p^{k_0}) := \{ (L_{i+p^{k_0}} \hookrightarrow V_{i+p^{k_0}} \twoheadrightarrow \mathcal{O}_X) \in \mathbb{E}_X \}_{i \in \mathbb{N}},$$

are k -points of $\mathcal{N}_{k_0} = \mathcal{N}_i$ for all $i \geq k_0$.

We define $T \subset \mathcal{N}_{k_0} \rightarrow S$ with its reduced structure to be the sub-scheme

$$T = \{ t \in \mathcal{N}_{k_0} \mid H^0(\mathcal{L}_t) = 0 \}.$$

It is open in \mathcal{N}_{k_0} . Let \mathcal{L} be the restriction of universal line bundle on $X_S \times_S T$ (thus, for any $t \in T$, $H^1(\mathcal{L}_t^{p^i}) \neq 0$ for $i \geq 0$). Unfortunately, $R^1 p_{T*}(\mathcal{L})$ is neither commuting with base change nor locally free in general, where $p_T : X_S \times_S T \rightarrow T$ is the projection. However,

$$\mathcal{E} = R^{n-1} p_{T*}(\mathcal{L}^\vee \otimes p_{X_S}^* \omega_{X_S/S}), \quad n = \dim(X)$$

may not be locally free, but does commute with base change since

$$H^n(\mathcal{L}_t^\vee \otimes \omega_{X_S \times \{t\}}) = H^0(\mathcal{L}_t)^\vee = 0, \quad \forall t \in T.$$

There exists a quotient scheme $\pi : M = \mathbb{P}(\mathcal{E}) \rightarrow T$ together with

$$\pi^* \mathcal{E} \rightarrow \mathcal{O}_M(1) \rightarrow 0,$$

which represents the functor that sends a T -scheme $p_W : W \rightarrow T$ to the set of isomorphic classes of quotients $p_W^* \mathcal{E} \rightarrow Q \rightarrow 0$, where Q is a line bundle on W ([7, Thm 2.2.4]). By definition, $M \xrightarrow{\pi} T \rightarrow S$ satisfies the properties (1) and (2) in the proposition (replacing M by M_{red} if necessary). The rest of the proof is devoted to the construction of a rational map

$$f : M \dashrightarrow M$$

over S , which satisfies the property (3) in the proposition.

Let $F : X_S \times_S T \rightarrow (X_S \times_S T)'$ denote the relative Frobenius morphism over T . We write

$$(3.5) \quad \begin{array}{ccccc} & & \xrightarrow{F_{X_S \times_S T}} & & \\ & \nearrow & & \searrow & \\ X_S \times_S T & \xrightarrow{F} & (X_S \times_S T)' & \longrightarrow & X_S \times_S T \\ & \searrow p_T & \downarrow p'_T & & \downarrow p_T \\ & & T & \xrightarrow{F_T} & T. \end{array}$$

One has $(X_S \times_S T)' = X'_S \times_S T$ thus we can project to X'_S , further to X_S . Let \mathcal{L}' be the pullback of \mathcal{L} under the projection

$$(X_S \times_S T)' \xrightarrow{p_1} X_S \times_S T$$

and ω (resp. ω') be the relative canonical line bundle of $X_S \times_S T \xrightarrow{p_T} T$ (resp. $(X_S \times_S T)' \xrightarrow{p'_T} T$). The line bundle $F^*\mathcal{L}'$ on $X_S \times_S T$ defines a S -morphism $\nu : T \rightarrow \text{Pic}_{X_S}^\tau$. Define

$$T^0 = \{t \in T \mid H^0(\mathcal{L}_t^p) = 0\} \subset T$$

with its reduced scheme structure. It is open in T . By definition of T , ν restricted to T^0 factors through T , thus induces

$$\nu : T^0 \rightarrow T$$

such that

$$(1 \times \nu)^*\mathcal{L} \otimes p_{T^0}^*\eta = F^*\mathcal{L}'$$

on $X_S \times_S T^0$, with $X_S \times_S T^0 \xrightarrow{p_{T^0}} T^0$, and where η is a line bundle on T^0 . We abuse notations and still write ω for the relative dualizing sheaf $\omega_{X_S \times_S T^0/T^0}$ of p_{T^0} , ω' for the relative dualizing sheaf $\omega_{X'_S \times_S T^0/T^0}$ of $p'_{T^0} : X'_S \times_S T^0 \rightarrow T^0$. Since $\omega = p_{X_S}^*\omega_{X_S/S} = (1 \times \nu)^*\omega_{X_S \times_S T/T}$, we have

$$R^{n-1}p_{T^0*}(\omega \otimes F^*\mathcal{L}'^\vee) = \eta^{-1} \otimes \nu^*\mathcal{E}, \quad R^{n-1}p'_{T^0*}(\omega' \otimes \mathcal{L}'^\vee) = F_{T^0}^*\mathcal{E},$$

and those identities commute with base change. By Lemma 3.6, we have

$$\phi : \eta^{-1} \otimes \nu^*\mathcal{E} \rightarrow F_{T^0}^*\mathcal{E}.$$

Together with the universal quotient $\pi^*\mathcal{E} \rightarrow \mathcal{O}_{M_0}(1) \rightarrow 0$, restricted to $M_0 := M \times_S T^0$, where $\pi : M_0 \rightarrow T^0$, $T^0 \rightarrow S$ factors through the composition of the open embedding $T^0 \subset T$ with the map $T \rightarrow S$, this induces

$$\Phi : \pi^*\nu^*\mathcal{E} \xrightarrow{\phi} \pi^*\eta \otimes \pi^*F_{T^0}^*\mathcal{E} = \pi^*\eta \otimes F_{M_0}^*\pi^*\mathcal{E} \rightarrow \pi^*\eta \otimes \mathcal{O}_{M_0}(p).$$

Here F_{T^0} , F_{M_0} are the absolute Frobenius morphisms satisfying

$$\begin{array}{ccc} M_0 & \xrightarrow{F_{M_0}} & M_0 \\ \downarrow \pi & & \downarrow \pi \\ T^0 & \xrightarrow{F_{T^0}} & T^0 \end{array}$$

Let $M^0 \subset M$ be the reduced open set consisting of points $q \in M$ such that Φ is surjective at $q \in M$ (which implies $M^0 \subset M_0$). Then there exists a unique morphism

$$f : M^0 \rightarrow M$$

corresponding via the universal property to $f^*(\pi^*\mathcal{E} \rightarrow \mathcal{O}_M(1) \rightarrow 0) = (\pi^*\nu^*\mathcal{E} \rightarrow \pi^*\eta \otimes \mathcal{O}_{M^0}(p) \rightarrow 0)$. By definition, one has the factorization

$$\begin{array}{ccc} M^0 & \xrightarrow{f} & M \\ \downarrow \pi & & \downarrow \pi \\ T^0 & \xrightarrow{\nu} & T. \end{array}$$

Then Lemma 3.6 below implies that the rational map

$$f : M \dashrightarrow M$$

satisfies the requirement (3) in the proposition. \square

For any point $t \rightarrow T$, let $s \rightarrow S$ be its image under $T \rightarrow S$, and X_s be the fiber of $X_S \rightarrow S$ at $s \rightarrow S$. Then the diagram (3.5) specializes to

$$\begin{array}{ccccc} & & F_{X_s \times t} & & \\ & \nearrow & & \searrow & \\ X_s \times t & \xrightarrow{F_t} & (X_s \times t)' & \longrightarrow & X_s \times t \\ & \searrow & \downarrow & & \downarrow \\ & & t & \longrightarrow & t \end{array}$$

and F_t induces the $k(t)$ -linear map $F_t^* : H^1(\mathcal{L}'_t) \rightarrow H^1((F^*\mathcal{L}')_t)$, which induces a $k(t)$ -linear map

$$(F_t^*)^\vee : H^1((F^*\mathcal{L}')_t)^\vee \rightarrow H^1(\mathcal{L}'_t)^\vee.$$

By Serre duality, we have the induced $k(t)$ -linear map

$$(3.6) \quad \phi_t : H^{n-1}((\omega \otimes F^*\mathcal{L}'^\vee)_t) \rightarrow H^{n-1}((\omega' \otimes \mathcal{L}'^\vee)_t).$$

Serre duality allows to make (3.6) in families:

Lemma 3.6. *There exists a homomorphism*

$$\phi : R^{n-1}p_{T*}(\omega \otimes F^*\mathcal{L}'^\vee) \rightarrow R^{n-1}p'_{T*}(\omega' \otimes \mathcal{L}'^\vee)$$

such that

$$\begin{array}{ccc} R^{n-1}p_{T*}(\omega \otimes F^*\mathcal{L}'^\vee) \otimes k(t) & \xrightarrow{\tau} & H^{n-1}((\omega \otimes F^*\mathcal{L}'^\vee)_t) \\ \downarrow \phi \otimes k(t) & & \downarrow \phi_t \\ R^{n-1}p'_{T*}(\omega' \otimes \mathcal{L}'^\vee) \otimes k(t) & \xrightarrow{\tau} & H^{n-1}((\omega' \otimes \mathcal{L}'^\vee)_t) \end{array}$$

is commutative for any $t \rightarrow T$, where τ is the canonical base change isomorphism, and ϕ_t is the homomorphism in (3.6).

Proof. Let ω (resp. ω') be the relative canonical line bundle of

$$X_T := X_S \times_S T \xrightarrow{p_T} T$$

(resp. $X'_T := (X_S \times_S T)' \xrightarrow{p'_T} T$). Let $F : X_T \rightarrow X'_T$ be the relative Frobenius morphism in (3.5), which is flat and Cohen-Macaulay as $X_S \rightarrow S$ is smooth. The injection $\mathcal{L}' \hookrightarrow F_*F^*\mathcal{L}'$ of vector bundles induces the surjection of vector bundles

$$(3.7) \quad \mathcal{H}om(F_*F^*\mathcal{L}', \omega') \twoheadrightarrow \mathcal{H}om(\mathcal{L}', \omega').$$

Duality theory for the Cohen-Macaulay map $F : X_T \rightarrow X'_T$ implies an isomorphism

$$(3.8) \quad F_*\mathcal{H}om(F^*\mathcal{L}', \omega) \cong \mathcal{H}om(F_*F^*\mathcal{L}', \omega').$$

Equations (3.7) and (3.8) imply that one has a surjection

$$(3.9) \quad F_*\mathcal{H}om(F^*\mathcal{L}', \omega) \twoheadrightarrow \mathcal{H}om(\mathcal{L}', \omega')$$

of vector bundles on X'_S . Since F is a finite morphism, taking $R^{n-1}p'_{T*}((3.9))$ induces

$$(3.10) \quad \phi : R^{n-1}p_{T*}\mathcal{H}om(F^*\mathcal{L}', \omega) \rightarrow R^{n-1}p'_{T*}\mathcal{H}om(\mathcal{L}', \omega').$$

For any $t \rightarrow T$, let X_t be the fiber of $X_T \rightarrow T$, and $F_t : X_t \rightarrow X'_t$ be the relative Frobenius. Then $(3.10) \otimes k(t)$ induces (through τ)

$$\phi \otimes k(t) : H^{n-1}(X_t, \mathcal{H}om(F_t^*\mathcal{L}'_t, \omega_t)) \rightarrow H^{n-1}(X'_t, \mathcal{H}om(\mathcal{L}'_t, \omega'_t))$$

which is induced by $\varphi := (3.9) \otimes k(t)$. Then the surjection of vector bundles on X'_t

$$(3.11) \quad \varphi : (F_t)_*\mathcal{H}om(F_t^*\mathcal{L}'_t, \omega_t) \twoheadrightarrow \mathcal{H}om(\mathcal{L}'_t, \omega'_t)$$

is dual, by taking $\mathcal{H}om(\cdot, \omega'_t)$, to the canonical injection

$$(3.12) \quad \mathcal{L}'_t \hookrightarrow (F_t)_*F_t^*\mathcal{L}'_t$$

of vector bundles on X'_t . As the dual $H^{n-1}(\varphi)^\vee$ of $H^{n-1}(\varphi) = \phi \otimes k(t)$ is the $k(t)$ -linear map

$$F_t^* : H^1(X'_t, \mathcal{L}'_t) \rightarrow H^1(X_t, F_t^* \mathcal{L}'_t)$$

induced by (3.12), this finishes the proof. \square

To prove Theorem 3.4, the key tool is a theorem of Hrushovski. In fact, we only need a special case of his theorem [6, Corollary 1.2].

Theorem 3.7 (Hrushovski, [6, Corollary 1.2]). *Let Y be an affine variety over \mathbb{F}_q , and let $\Gamma \subset (Y \times_{\mathbb{F}_q} Y) \otimes_{\mathbb{F}_q} \overline{\mathbb{F}}_q$ be an irreducible subvariety over $\overline{\mathbb{F}}_q$. Assume the two projections $\Gamma \rightarrow Y$ are dominant. Then, for any closed subvariety $W \subsetneq Y$, there exists $x \in Y(\overline{\mathbb{F}}_q)$ such that $(x, x^{q^m}) \in \Gamma$ and $x \notin W$ for large enough natural number m .*

Remark 3.8. Writing $Y \subset \mathbb{A}^r$, then $x = (x_1, \dots, x_r) \in (\overline{\mathbb{F}}_q)^r = \mathbb{A}^r(\overline{\mathbb{F}}_q)$ and $x^{q^m} := (x_1^{q^m}, \dots, x_r^{q^m}) \in \mathbb{A}^r(\overline{\mathbb{F}}_q)$. In our application, Γ is the Zariski closure of the graph of a dominant rational map $f : Y \dashrightarrow Y$ and W is the locus where f is not well-defined.

Proof of Theorem 3.4. (Compare with [3, Thm 3.14].) Let

$$f : M \dashrightarrow M$$

be as in Proposition 3.5. So $M \rightarrow S$ is a S -reduced scheme of finite type, where S is smooth affine over \mathbb{F}_q , the normalization of \mathbb{F}_p in $H^0(S, \mathcal{O}_S)$. Let M_k be the general fiber of $M \rightarrow S$ at $\text{Spec}(k) \rightarrow S$. We define Z_k to be the intersection of the Zariski closures $\overline{\mathbb{E}_X(m)}$ of the $\mathbb{E}_X(m) \subset M_k$, where $\mathbb{E}_X(m) = \{ (L_{i+m} \hookrightarrow V_{i+m} \twoheadrightarrow \mathcal{O}_X) \in \mathbb{E}_X \}_{i \in \mathbb{N}}$. By the noetherian property, there exists a $m_0 > 0$ such that

$$Z_k = \overline{\mathbb{E}_X(m)} \subset M_k(k) = M(k)$$

for any $m \geq m_0$. As \mathbb{E}_X is assumed to be infinite, all components of Z_k have dimension ≥ 1 . Indeed, if it had a component of dimension 0, then there would be a $m_1 > m_0$ such that this 0-dimensional component would not lie on $\mathbb{E}_X(m_1)$, a contradiction. The fields of constants of the irreducible components of Z_k lies between $\mathbb{F}_q(S)$ and k . Let $S' \rightarrow S$ be affine with S' smooth affine irreducible such that all components of Z_k are defined over S' , yielding models of them over S' which are geometrically irreducible over \mathbb{F}_q , and a model $Z \rightarrow S'$ of Z_k . We replace now $X_S \rightarrow S$ and $M \rightarrow S$ by $X_{S'} \rightarrow S'$ and the corresponding $M_{S'} \rightarrow S'$. We abuse notations, set $S = S'$, and have $Z \subset M \rightarrow S$.

Let $M^0 \subset M$ be the largest open subset where f is well-defined. Notice that

$$\mathbb{E}_X(1) \subset M_k^0$$

and $f(\mathbb{E}_X(m)) = \mathbb{E}_X(m-1)$ for any $m \geq 1$. Thus $f(Z_k \cap M_k^0)$ contains a dense subset of Z_k . On the other hand,

$$f(Z_k \cap M_k^0) \subset Z_k,$$

thus $f : M \dashrightarrow M$ induces a dominant rational map $f : Z_k \dashrightarrow Z_k$, and thus a rational dominant map

$$f : Z \dashrightarrow Z.$$

We consider one irreducible component of Z_k , and its model $Z^{\text{irr}} \rightarrow S$, which is a geometrically irreducible component of Z over \mathbb{F}_q . There is an integer $a_1 > 0$ such that $f^{a_1} : Z^{\text{irr}} \dashrightarrow Z^{\text{irr}}$ is a dominant rational map over a finite field \mathbb{F}_q .

To prove that Z^{irr} contains a periodic point of f^{a_1} , without loss of generality, we can assume that $Y := Z^{\text{irr}} \subset \mathbb{A}^r$ is an affine variety and $f^{a_1} : Y \dashrightarrow Y$ is defined by rational functions $f_1, \dots, f_r \in \mathbb{F}_q(Y)$. Let

$$\Gamma \subset Y \times_{\mathbb{F}_q} Y$$

be the Zariski closure of the graph of f^{a_1} , which is irreducible over $\overline{\mathbb{F}}_q$ as Y is by assumption. The two projections $\Gamma \rightarrow Y$ are dominant since $f^{a_1} : Y \dashrightarrow Y$ is dominant. By Theorem 3.7, there exists a point $x \in Y(\overline{\mathbb{F}}_q)$ such that f is well-defined at $x = (x_1, \dots, x_r)$ and $f^{a_1}(x) = (x_1^{q^m}, \dots, x_r^{q^m}) := x^{q^m}$ for some large m . Recall that $f_i \in \mathbb{F}_q(Y)$ ($i = 1, \dots, r$) have coefficients in \mathbb{F}_q , we have

$$\begin{aligned} f^{a_1}(f^{a_1}(x)) &= (f_1(x_1^{q^m}, \dots, x_r^{q^m}), \dots, f_r(x_1^{q^m}, \dots, x_r^{q^m})) \\ &= (f_1(x_1, \dots, x_r)^{q^m}, \dots, f_r(x_1, \dots, x_r)^{q^m}) \\ &= (x_1^{q^{2m}}, \dots, x_r^{q^{2m}}) = x^{q^{2m}}. \end{aligned}$$

Thus $f^{a_2 a_1}(x) = x^{q^{a_2 a_1}} = x$ when a_2 is large enough. The point

$$x \in Y(\overline{\mathbb{F}}_q)$$

determines a nontrivial extension $0 \rightarrow L \rightarrow V \rightarrow \mathcal{O}_{X_{\bar{s}}} \rightarrow 0$ on a good reduction $X_{\bar{s}}$ of X (over $\overline{\mathbb{F}}_q$). That $f^a(x) = x$, $a = a_1 a_2$, means by Proposition 3.5 (3)

$$((F_{X_{\bar{s}}}^*)^a L \hookrightarrow (F_{X_{\bar{s}}}^*)^a V \twoheadrightarrow (F_{X_{\bar{s}}}^*)^* \mathcal{O}_{X_{\bar{s}}}) \cong (L \hookrightarrow V \twoheadrightarrow \mathcal{O}_{X_{\bar{s}}})$$

up to scale. This finishes the proof of Theorem 3.4 and thus of Theorem 3.1. \square

Theorem 3.9. *Let X be a smooth projective variety over an algebraically closed field k of characteristic $p > 0$, then:*

- (i) *Every stratified bundle on X is trivial if and only if π_1 is trivial.*

- (ii) *All the irreducible stratified bundles have rank 1 if and only if $[\pi_1, \pi_1]$ is a pro- p -group.*
- (iii) *Every stratified bundle is a direct sum of stratified line bundles, that is $\mathbf{str}(X)$ is a semi-simple category with irreducible objects of rank 1, if and only if π_1 is abelian with no non-trivial p -power quotient.*

Proof. (i) is the main theorem of [3]. (ii) is Theorem 2.6. To show (iii), assume that π_1 is abelian with no p -power order quotient. Let $E = (E_n, \sigma_n)_{n \in \mathbb{N}}$ be a stratified bundle on X . By (ii), any irreducible stratified bundle has rank 1. Thus there is a filtration $0 = E^0 \subset E^1 \subset \dots \subset E^r = E$ in $\mathbf{str}(X)$ such that $L^v = E^v/E^{v-1}$, $1 \leq v \leq r$, are rank one stratified bundles. Then, by Theorem 3.1, the filtration splits and E is a direct sum of rank 1 objects. \square

We now comment on analogs of (ii) and (iii) in complex geometry. For this reason, we include here the following lemma, which may have an independent interest.

Lemma 3.10. *Let G be a commutative group scheme over an algebraically closed field k such that all its quotients in \mathbb{G}_m are smooth. Let ℓ be a non-trivial character of G . Then $H^1(G, \ell) = 0$.*

Proof. Let $\chi : G \rightarrow \mathbb{G}_m = \text{Aut}(\ell)$ be the non-trivial character, and let $\sigma : G \rightarrow \ell$ be a cocycle representing a class in $H^1(G, \ell)$. By definition of a cocycle, one has $\sigma(gh) = \chi(g)\sigma(h) + \sigma(g)$. The commutativity of G implies

$$(3.13) \quad \sigma(hg) = \chi(h)\sigma(g) + \sigma(h) = \sigma(gh) = \chi(g)\sigma(h) + \sigma(g).$$

As χ is non-trivial, and $\text{Im}(\chi) \subset \mathbb{G}_m$ is assumed to be smooth, there is a $h \in G(k)$ such that

$$0 \neq (\chi(h) - 1) \in \text{End}(\ell) = k.$$

Thus

$$(\chi(h) - 1) \in k^\times = \mathbb{G}_m(k) = \text{Aut}(\ell).$$

Set $v = (\chi(h) - 1)^{-1}\sigma(h) \in \ell$. Then, by (3.2), one has $\sigma(g) = \chi(g)v - v$, which means that σ is a coboundary. Thus $H^1(G, \ell) = 0$. \square

Remark 3.11. The same proof shows that if X is a smooth complex variety, with abelian topological fundamental group, and if ℓ is a non-trivial rank 1 local system, then $H^1(X, \ell) = 0$. Indeed, G is now $\pi_1^{\text{top}}(X, a)$, the topological fundamental group based at a complex point $a \in X(\mathbb{C})$, and non-triviality implies the existence of $h \in G$ with $(\chi(h) - 1) \in \mathbb{C}^\times = \text{Aut}(\ell)$. One then concludes identically.

This fact ought to be well known, but we could not find a reference in the literature.

Remark 3.12. As already mentioned in the introduction, the Malcev-Grothendieck theorem ([10], [5]) asserts that the étale fundamental group π_1 of a smooth complex projective variety is trivial if and only if its stratified fundamental group π^{str} is trivial, where π^{str} is the pro-affine complex algebraic group of the Zariski closures of the monodromies of complex linear representations of the topological fundamental group π_1^{top} . By going to the associated Galois cover, it implies that π^{str} is finite if and only if π_1 is finite (in which case $\pi^{\text{str}} = \pi_1$). One also has that π^{str} is abelian if and only if π_1 is abelian. Indeed, by definition π^{str} is abelian if and only if irreducible complex linear representations have dimension 1. But if $\rho : \pi_1^{\text{top}} \rightarrow GL(r, \mathbb{C})$ is a representation, since π_1^{top} is spanned by finitely many elements, ρ has values in $GL(r, A)$ for A a ring of finite type over \mathbb{Z} . Then if ρ is irreducible, there is a closed point $s \in \text{Spec}(A)$ such that $\rho \otimes \kappa(s) : \pi_1^{\text{top}} \rightarrow GL(r, \kappa(s))$ is irreducible as well, where $\kappa(s)$ is the residue field of s . As $\rho \otimes \kappa(s)$ is finite, it factors through π_1 , thus $r = 1$. Since a linear representation with finite monodromy is semi-simple, one can summarize as follows:

Claim 3.13. π_1 is

- 1) finite,
- 2) resp. abelian,
- 3) resp. abelian and finite

if and only if π_1^{str}

- 1) finite
- 2) resp. abelian,
- 3) resp. abelian and finite

if and only if

- 1) local systems have finite monodromy,
- 2) resp. irreducible local systems have rank 1,
- 3) resp. local systems are direct sums of rank 1 local systems.

While the main result of [3] is an analog in characteristic $p > 0$ of Claim 3.13 1), the main theorem of this note is an analog of Claim 3.13 2) and 3).

REFERENCES

- [1] B. Conrad: *Grothendieck Duality and Base Change*, Lecture Notes in Mathematics **1750**, 2000.
- [2] J.P.P. dos Santos: *Fundamental group schemes for stratified sheaves*, Journal of Algebra, **317**, (2007), 691-713.

- [3] H. Esnault and V. Mehta: *Simply connected projective manifolds in characteristic $p > 0$ have no nontrivial stratified bundles*, Invent. math. **181** (2010), 449–465.
- [4] D. Gieseker: *Flat vector bundles and the fundamental group in non-zero characteristics*, Ann. Sc. Norm. Super. Pisa, **4** Ser. 2 (1), (1975), 1–31.
- [5] A. Grothendieck: *Représentations linéaires et compactifications profinies des groupes discrets*, Manuscripta mathematica, **2** (1970), 375–396.
- [6] E. Hrushovski: *The Elementary Theory of the Frobenius Automorphism*, arXiv: math/0406514v1 [math.LO], 2004.
- [7] D. Huybrechts, M. Lehn: *The Geometry of the Moduli Space of Sheaves*, Aspects of Mathematics **E31** Vieweg Verlag, Braunschweig 1997.
- [8] S. Kleiman: *The Picard Scheme*, Fundamental Algebraic Geometry, in Mathematical Surveys and Monographs, vol **123** (2005), 235–300.
- [9] H. Lange and U. Stuhler: *Vektorbündel auf Kurven und Darstellungen der algebraischen Fundamentalgruppe*, Math. Z., **156**, (1977), 73–83.
- [10] A. Malcev: *On isomorphic matrix representations of infinite groups*, Rec. Math. [Mat. Sbornik] N.S. **8** (50) (1940), 405–422.
- [11] D. Mumford: *Abelian Varieties*, Oxford University Press, 1970.
- [12] J.-P. Serre: *Linear Representations of Finite Groups*, Graduate Texts in Mathematics **42**, 1977.
- [13] SGA1.: *Revêtements étales et groupe fondamental*, SGA 1.

MATHEMATIK, UNIVERSITÄT DUISBURG-ESSEN, 45117 ESSEN, GERMANY
E-mail address: esnault@uni-due.de

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCE, BEIJING, P. R. OF CHINA
E-mail address: xsun@math.ac.cn